

# Differential effects of endurance, interval, and resistance training on telomerase activity and telomere length in a randomized, controlled study

Christian M. Werner<sup>1\*†</sup>, Anne Hecksteden<sup>2†</sup>, Arne Morsch<sup>3</sup>, Joachim Zundler<sup>1</sup>,  
Melissa Wegmann<sup>2</sup>, Jürgen Kratzsch<sup>4</sup>, Joachim Thiery<sup>4</sup>, Mathias Hohl<sup>1</sup>,  
Jörg Thomas Bittenbring<sup>5</sup>, Frank Neumann<sup>5</sup>, Michael Böhm<sup>1</sup>,  
Tim Meyer<sup>2‡</sup>, and Ulrich Laufs<sup>6‡</sup>

<sup>1</sup>Klinik für Innere Medizin III, Kardiologie, Angiologie und Internistische Intensivmedizin, Universität und Universitätsklinikum des Saarlandes, Geb. 41.1/IMED, Homburg/Saar 66421, Germany; <sup>2</sup>Institut für Sport und Präventivmedizin, Universität des Saarlandes, Campus, B8 2, Saarbrücken 66123, Germany; <sup>3</sup>Deutsche Hochschule für Prävention und Gesundheitsmanagement, Hermann-Neuberger-Sportschule 3, Saarbrücken 66123, Germany; <sup>4</sup>Institut für Labormedizin, Klinische Chemie und Molekulare Diagnostik, Universitätsklinikum, Liebigstr. 20, Leipzig 04103, Germany; <sup>5</sup>Klinik für Innere Medizin I, Onkologie, Hämatologie, Klinische Immunologie und Rheumatologie, Universitätsklinikum des Saarlandes, Geb. 41.1/IMED, Homburg/Saar 66421, Germany; and <sup>6</sup>Klinik und Poliklinik für Kardiologie, Universitätsklinikum Leipzig, Liebigstr. 20, 04103 Leipzig, Germany

Received 11 November 2017; revised 26 February 2018; editorial decision 23 August 2018; accepted 21 September 2018

## Aims

It is unknown whether different training modalities exert differential cellular effects. Telomeres and telomere-associated proteins play a major role in cellular aging with implications for global health. This prospective training study examines the effects of endurance training, interval training (IT), and resistance training (RT) on telomerase activity and telomere length (TL).

## Methods and results

One hundred and twenty-four healthy previously inactive individuals completed the 6 months study. Participants were randomized to three different interventions or the control condition (no change in lifestyle): aerobic endurance training (AET, continuous running), high-intensive IT (4 × 4 method), or RT (circle training on 8 devices), each intervention consisting of three 45 min training sessions per week. Maximum oxygen uptake (VO<sub>2max</sub>) was increased by all three training modalities. Telomerase activity in blood mononuclear cells was up-regulated by two- to three-fold in both endurance exercise groups (AET, IT), but not with RT. In parallel, lymphocyte, granulocyte, and leucocyte TL increased in the endurance-trained groups but not in the RT group. Magnet-activated cell sorting with telomerase repeat-amplification protocol (MACS-TRAP) assays revealed that a single bout of endurance training—but not RT—acutely increased telomerase activity in CD14+ and in CD34+ leucocytes.

## Conclusion

This randomized controlled trial shows that endurance training, IT, and RT protocols induce specific cellular pathways in circulating leucocytes. Endurance training and IT, but not RT, increased telomerase activity and TL which are important for cellular senescence, regenerative capacity, and thus, healthy aging.

## Keywords

Exercise • Cellular senescence • Telomere length • Telomerase activity • Endurance training  
• Resistance training

\* Corresponding author. Tel/Fax: +49 6841 161 5000/15175, Email: christian.werner@uks.eu

† The first two authors contributed equally to this manuscript as first authors.

‡ The last two authors contributed equally to this manuscript as senior authors.

© The Author(s) 2018. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

## Translational perspective

Exercise is a cornerstone of cardiovascular disease prevention. This randomized, controlled, and prospective training study shows that specific training protocols lead to differential effects on cellular aging. Aerobic endurance and high-intensive interval training, but not resistance training, increases telomerase activity and telomere length in blood mononuclear cells. The study identifies telomerase activity and telomere length as sensitive cellular parameters to measure preventive effects of exercise interventions. Individual training recommendations guided by measurement of parameters associated with cellular senescence and training response may improve both adherence to and efficacy of exercise training programmes in cardiovascular prevention.

## Introduction

Physical training is associated with improvements in many aspects of human health. These include exercise capacity, endothelial function, insulin sensitivity, autonomic function, and blood pressure as well as a reduction of abdominal fat, lipid metabolism, adverse psychosocial, and inflammatory parameters.<sup>1,2</sup> Endurance training and resistance training (RT) alone or in combination are recommended for cardiovascular disease prevention.<sup>2–4</sup> Compared with conventional aerobic endurance training (AET), high-intensity interval training (IT) is believed to be a more effective and time-efficient training method for improving cardiorespiratory fitness.<sup>5</sup> However, few randomized comparisons of different training modalities have been reported, and it is unknown whether they exert differential cellular effects.

Vascular aging is associated with endothelial dysfunction and atherogenesis.<sup>6</sup> A key cellular process of aging is the shortening of telomeres to a critical telomere length (TL) where cells enter replicative senescence or programmed cell death.<sup>7</sup> Telomere length in circulating leucocytes is associated with cardiovascular morbidity.<sup>6,8–11</sup> Telomere length is regulated by the reverse transcriptase telomerase that prevents successive shortening of telomeres.<sup>7,12</sup> The regulation of telomerase activity and TL in blood mononuclear cells (MNCs) parallels the regulation in the vessel wall and telomeres shorten at similar rates in somatic tissues of adults.<sup>8,10,11,13,14</sup>

Beneficial effects of physical activity on cellular regeneration and senescence have been observed.<sup>10,12,13,15–19</sup> For example, long-term endurance training is associated with higher telomerase activity and a reduced rate of telomere attrition in young- and middle-aged endurance athletes compared with inactive controls.<sup>13</sup> Twins with a higher level of physical activity exhibit longer telomeres in middle age compared with inactive siblings.<sup>12</sup> Moderate levels of physical activity may reduce telomere shortening by inducing a telomere-protective phenotype, indicating an important cellular adaptation that may prevent age-related diseases.<sup>18</sup> However, prospective studies are lacking, and the effect of different exercise modalities is unknown.<sup>12</sup>

Herein, we compare the long-term effects of three exercise modes, namely AET, high-intensity IT, and RT, with a sedentary control group on telomerase activity and TL in a prospective, randomized study in a primary prevention cohort. In addition, we measured the acute effects of endurance training and RT on telomerase activity in leucocytes subpopulations in a cross-over study.

## Methods

### Acute effects of aerobic endurance and resistance exercise

Exercise testing was performed by bicycle spiroergometry (Innocor 500; Innovision, Denmark; 30 W/min ramp protocol) in young healthy, moderately trained individuals. The acute exercise study consisted of two supervised exercise bouts of running the maximum distance for 45 min at separate days in a cross-over design,  $n = 15$ . Resistance exercise was circle training on eight strength training devices (details in [Supplementary material online, Methods](#)).

### Prospective training study

The flowchart of the trial is illustrated in [Figure 2](#). In total, 1534 volunteers were screened. Two hundred and sixty-six healthy, physically inactive non-smokers without permanent medication were randomized. The main inclusion criteria were selected to represent the typical beginner of a primary preventive training programme: age 30–60 years, a body mass index (BMI) below 30 kg/m<sup>2</sup>, and a lack of regular physical exercise (<1 h/week in the last 12 months; VO<sub>2max</sub> <45 mL/kg/min). The exclusion criteria were smoking, known cardiovascular disease, cardiovascular risk factors requiring medication (including fasting blood glucose concentration <6.1 mmol/L, resting blood pressure >160/100 mmHg, and presence of iron deficiency/anaemia), any condition or disease that may interfere with the training protocol. Subjects were recruited by advertisements in local media. Sex and VO<sub>2max</sub> were used as criteria for stratified randomization with three strata for VO<sub>2max</sub> (<30 mL/kg/min; 30–40 mL/kg/min and >40 mL/kg/min). During the study, 119 subjects dropped out, primarily because of lack of compliance with the training protocol. The rate of drop-outs and injuries was higher in the two endurance groups. Twenty-three individuals had to be excluded due to lack of exhaustion on the stress tests or missing samples.

### Exercise testing

Exhaustive individual exercise tests were carried out using a combined step-wise and ramp-wise protocol as described in detail in the [Supplementary material online, Methods](#).<sup>20</sup>

### Training interventions

The details are described in the [Supplementary material online, Methods](#). Subjects in the control group were advised to maintain their lifestyle and diet for the next 6 months. Subjects in the training groups exercised three times per week for 26 weeks. Aerobic endurance training consisted of 45 min of walking/running at 60% heart rate reserve. Interval training was performed according to the high-intensity 4 × 4 method. Resistance training was a circle training of eight machine-based exercises: back extension, crunch, pull-down, seated row, seated leg curl, seated leg extension, seated chest press, and lying leg press. Twenty repetition

maximum (RM) was determined every 6 weeks and training weights were adjusted.

### Isolation of mononuclear cells

Mononuclear cells were isolated by Ficoll density gradient centrifugation before, after, and after 24 h in the acute exercise study.<sup>13</sup> In the prospective exercise study, MNCs were isolated at baseline and 48 h to 7 days after that last exercise bout after completion of the 6 months training intervention or control condition (see [Supplementary material online, Methods](#) for details).

### Telomerase activity

Telomerase activity was quantified using the Telomerase Repeat Amplification Protocol<sup>10,13,16,21</sup> in protein extracts of 10 000 MNCs or MACS-isolated leucocyte subpopulations using a Lightcycler (Roche, Germany). Protein extracts from human embryonic kidney (HEK) cells were measured as positive controls in each assay, a 10-point standard titration curve was established from 5 to 2500 cells to ensure linearity ([Supplementary material online, Figure S2](#),  $R^2 = 0.99$ ). Details are described in the [Supplementary material online, Methods](#).

### Telomere length

Telomere length in MNCs and different leucocyte subpopulations was determined by three separate methods, namely flow cytometry and FISH (FlowFISH) as well as by real-time PCR as described. Details are given in the [Supplementary material online, Methods](#). Southern blots were performed from the genomic DNA to standardize conversion of fluorescence values to base pairs.

### PCR method

Briefly, 40 ng of genomic DNA were used for real-time PCR using 150 nmol/L primers for the single copy gene 36b4 (S) and telomere strands (T).<sup>8,13,22</sup> Experiments were run on a StepOnePlus Real-Time PCR System (Applied Biosystems). The comparative cycle threshold (Ct) method ( $2^{-\Delta\Delta C_t}$ ) was used to calculate telomere/single copy gene (T/S) ratios.

### FlowFISH method

A telomere-specific green fluorescent peptide nucleic acid (PNA) probe (FITC-PNA, Panagene, Korea, via Cambridge Research Biochemicals, UK; sequence: FITC-OO-CCCTAA-CCCTAA-CCCTAA) was used to specifically bind telomeres.<sup>13,23</sup> Telomere fluorescence intensity was measured in different blood cell populations, namely lymphocytes and granulocytes as well as in bovine thymocytes from a single isolation, which were used as an internal standard in each sample.<sup>23</sup> Only MNCs with effective DNA counterstaining and effective PNA hybridization were gated. Lymphocytes and granulocytes were differentiated from thymocytes by scatter graphs, DNA signal, and gating in the FITC channel. From fluorescence intensities, molecules of soluble fluorescence (MESF) were calculated using calibration beads with defined fluorescence intensity. Additional validation experiments are listed in the [Supplementary material online, Methods](#).

### Southern blot determination of terminal restriction fragment length

The detailed procedure using the TeloTAGGG Telomere Length Assay Kit is described in the [Supplementary material online, Methods](#). Terminal restriction fragment values of MNC samples of 20 participants of the chronic exercise study were plotted against their corresponding FlowFISH-based MESF values to generate a standard curve ([Supplementary material online, Figure S4](#)), which was used to calculate the TLs for the other samples.

### Fluorescence-activated cell sorting (FACS) assay of leucocyte subpopulations in acute and chronic exercise

Characterization of leucocyte subpopulations was performed by multi-colour flow cytometry (see [Supplementary material online, Methods](#)).

### Telomerase activity in magnetic bead-isolated leucocyte subpopulations

CD34+ haematopoietic stem/progenitor cells as well as defined subpopulations of peripheral MNCs were isolated by magnetic-activated cell sorting (MACS) with magnetic bead-coated antibodies (Miltenyi Biotec MicroBeads; CD34: 130-046-703; CD4: 130-045-101; CD8: 130-045-201; CD14: 130-050-201; CD19: 130-050-301; CD56: 130-050-401) followed by magnetic enrichment and TRAP assay. A scheme of the isolation is provided in [Supplementary material online, Figure S6](#). Tests of purity and viability are provided in [Supplementary material online, Figure S7](#).

### Quantification of cortisol and serum markers of exercise

Analysis of serum cortisol was performed by liquid chromatography tandem mass spectrometry (LC-MS/MS) as described. As control for extreme physical activity, samples from  $n = 10$  participants of the 38th Berlin Marathon were included.<sup>24</sup> For more details see the [Supplementary material online, Methods](#).

### Statistical methods

Data are presented as individual data points with mean (bars) and standard deviation (SD, whiskers). For the two acute exercise cross-over studies, changes of telomerase activity over time were analysed by one-way analysis of variance (ANOVA) for repeated measurements within each training modality (*post hoc* test for linear trend). In the randomized controlled study, baseline characteristics between groups were compared by one-way ANOVA. Changes in blood composition, serum markers, chemistry, clinical variables, and physical fitness during the study period were analysed using *t*-tests for related samples and two-way ANOVA for repeated measurements (repeat factor: test; categorical predictor: group; *post hoc* test: Bonferroni's multiple comparison test) to compare groups. Changes of TL and telomerase activity within groups were analysed with *t*-tests and differences between groups were tested with one-way ANOVA and the Bonferroni *post hoc* tests for multiple comparisons. Generalized estimating equation (GEE) mixed models were employed to test the influence of study group (dummy coded), age, gender, BMI, and baseline  $VO_{2max}$  on the changes TL and telomerase activity from pre- to post-test. Missing samples were excluded. Kolmogorov-Smirnov tests were used to test the hypothesis of normal distribution and all variables used in GEE models were normally distributed. Associations between molecular regulators, clinical variables, and parameters of exercise capacity were tested by two-sided Pearson correlation. Differences in telomerase activity stratified by training response were analysed by *t*-tests. Analyses were performed with SPSS 20.0 (IBM Corporation, Armonk, NY, USA) and GraphPad Prism 5.01 (San Diego, CA, USA). *P*-values <0.05 were considered significant.

### Study approval

The clinical study has been conducted according to the principles of Helsinki and was approved after institutional review by the ethics committee of the Ärztekammer des Saarlandes (No. 148/10) and registered at [clinicaltrials.gov](#) (NCT01263522). Written informed consent was received from all participants prior to any study-related procedure. Names of study subjects were blinded by pseudonymization.

**Table 1** Baseline characteristics in the prospective 6 months training study

	Control group	Aerobic endurance training	Interval training	Resistance training	P-value
N = 124	35	26	29	34	
Gender (M/F)	12/23	9/17	10/19	14/20	
Age (years)	50.2 (7.4)	49.5 (7.0)	48.4 (6.5)	48.1 (7.5)	0.61
Clinical characteristics					
Body mass index (kg/m <sup>2</sup> )	24.2 (3.1)	23.8 (3.3)	24.5 (3.0)	24.8 (3.4)	0.64
Systolic BP (mmHg)	131 (16)	132 (15)	125 (27)	132 (13)	0.35
Diastolic BP (mmHg)	84 (10)	86 (11)	82 (17)	86 (7)	0.63
Resting heart rate (b.p.m.)	64 (10)	65 (8)	68 (11)	67 (9)	0.32
Clinical chemistry					
Fasting glucose (mg/dL)	96 (9)	96 (6)	98 (8)	96 (8)	0.44
Fasting insulin (μU/mL)	4.60 (2.2)	4.48 (2.3)	6.11 (2.6)	5.19 (2.6)	0.05
Total cholesterol (mg/dL)	212 (40)	224 (43)	212 (36)	206 (44)	0.41
HDL cholesterol (mg/dL)	55 (20)	58 (21)	53 (17)	56 (16)	0.77
LDL cholesterol (mg/dL)	105 (31)	114 (36)	106 (32)	102 (32)	0.59
Treadmill stress test					
Peak heart rate (b.p.m.)	178 (13)	183 (10)	178 (10)	182 (11)	0.19
VO <sub>2max</sub> (mL/min/kg)	35.1 (5.3)	35.3 (6.3)	35.1 (5.0)	35.3 (5.3)	0.99
Maximum speed (km/h)	10.0 (1.5)	9.9 (1.6)	10.0 (1.2)	10.0 (1.5)	0.99
At 130 b.p.m.	5.5 (2.4)	4.5 (3.1)	4.5 (2.8)	5.6 (1.9)	0.14
At 150 b.p.m.	7.2 (1.3)	7.2 (0.9)	6.9 (0.6)	7.2 (1.1)	0.56
At lactate 2 mmol/L	4.9 (3.0)	4.6 (3.2)	4.7 (3.0)	5.3 (2.6)	0.82
Respiratory exchange rate	1.17 (0.10)	1.12 (0.09)	1.10 (0.13)	1.11 (0.22)	0.28

Data are represented as mean (SD).

BP, blood pressure; b.p.m., beats per minute; F, female; M, male; VO<sub>2max</sub>, maximum oxygen uptake.

## Results

### Differential regulation of telomerase activity in mononuclear cell and MACS-isolated leucocyte subpopulations by acute exercise


The acute effects of aerobic endurance vs. resistance exercise were studied in  $N = 15$  healthy volunteers in a cross-over design (Supplementary material online, Table S1). All individuals performed supervised cardiopulmonary exercise testing until exertion (mean respiratory coefficient 1.1), which showed no cardiac or pulmonary limitation and a moderately trained status (VO<sub>2max</sub>  $41.4 \pm 6.3$  mL/min/kg). Volunteers ran a mean distance of  $7.3 \pm 1.5$  km and had a mean exercise heart rate of  $155 \pm 8$  b.p.m. and a peak exercise heart rate of  $178 \pm 7$  b.p.m., equalling a high-endurance intensity of 0.83 as calculated by the Karvonen formula. In the resistance exercise group, a mean weight of  $10.3 \pm 4.2$  tons was moved.

Aerobic endurance training acutely increased telomerase activity (pre:  $189 \pm 90$ , post:  $250 \pm 140$ , and 24 h:  $306 \pm 182$  HEK cell equivalents) (Figure 1A). This regulation was not observed when the same individuals performed intensive RT for the same duration. As depicted in Supplementary material online, Table S2, the endurance protocol increased absolute numbers of leucocytes, neutrophilic, and basophile granulocytes and monocytes. After 24 h, the blood counts returned to the baseline levels. Compared with these profound acute

changes of blood composition after the endurance exercise bout, changes were much smaller after resistance exercise. Fluorescence-activated cell sorting (FACS) analyses showed that different T cell populations (T-helper, T-killer, and activated T cells) were increased and there was a numeric trend of increased B and NK cells. The strongest up-regulation was seen in CD34+ haematopoietic stem/progenitor cells (Supplementary material online, Table S3). However, this pattern was not observed after resistance exercise. CD14+ monocytes and CD34+ progenitor cells exhibited a potent increase of telomerase activity directly and 24 h after endurance exercise (four-fold in CD14+ and six-fold in CD34+ cells, Figure 1B and C), whereas RT was not associated with these changes. Telomerase activity was not detectable in CD4+, CD8+, CD19+, and CD56+ cells.

### Prospective training study

The design of the randomized controlled training study is illustrated in Figure 2. The training groups were similar with respect to clinical characteristics, glucose and lipid parameters, and physical fitness. Table 1 shows the baseline characteristics. The maximum oxygen uptake (VO<sub>2max</sub>) was  $35.2 \pm 5.4$  mL/min/kg. The 6 months training interventions improved physical fitness. All three exercise modalities led to a comparable increase of the  $\Delta$ VO<sub>2max</sub> (in mL/min/kg: control  $-1.0 \pm 3.1$ ; AET  $2.7 \pm 3.7$ ; IT  $2.8 \pm 5.1$ ; RT  $3.0 \pm 5.9$ ). Maximum running speed as an indicator of maximum physical capacity was increased in



**Figure 1** Acute effects of aerobic endurance vs. resistance exercise on telomerase activity. (A) Cross-over comparison of acute regulation of mononuclear cell telomerase activity pre, post, and 24 h after aerobic endurance training (45 min continuous running) and resistance exercise (45 min circle training on eight strength devices) in  $n = 15$  healthy young individuals. Exercise was performed at least 48 h after any previous physical exercise. Telomerase activity of  $10^4$  mononuclear cell was compared with human embryonic kidney cells (telomere repeat amplification protocols, TRAP assay). Time course of individual % changes of telomerase activity in  $10^4$  magnetic-activated cell sorting-isolated (B) CD14+ and (C) CD34+ leukocyte subfractions compared with pre-exercise in  $n = 10$  healthy young individuals (blue circles: endurance exercise; green circles: resistance exercise).

all training groups. All training modalities reduced resting heart rate. Data depicted in [Table 2](#) and [Supplementary material online, Table S4](#).

### Differential regulation of leucocyte telomere length by endurance and resistance training

Telomere length was measured by FACS-FlowFISH and real-time PCR. After 6 months, a small decline of TL was observed in the

control group and in the RT group. In contrast, the two endurance-based training groups, the AET, and the IT, showed a significant increase of both lymphocyte and granulocyte TL (AET:  $\Delta$  lymphocyte TL  $218 \pm 211$  bp,  $\Delta$  granulocyte TL  $248 \pm 349$  bp; IT:  $\Delta$  lymphocyte TL  $214 \pm 307$  bp,  $\Delta$  granulocyte TL  $261 \pm 332$  bp; [Figure 3A](#) and [B](#)). Comparisons of the relative intra-individual changes of TL yielded a similar result with a non-significant decline of TL in the control and RT groups, while lymphocytes and granulocytes in the AET and IT groups were characterized by 3.3–3.5% longer telomeres ([Figure 3D](#)



**Table 2** Effects of 6 months training on clinical parameters and physical fitness

	Co	Aerobic endurance training		Interval training		Resistance training	
	Δ post vs. pre	Δ post vs. pre	P vs. Co	Δ post vs. pre	P vs. Co	Δ post vs. pre	P vs. Co
Clinical parameters							
BMI (kg/m <sup>2</sup> )	0.0 (0.8)	-0.5 (1.0)	<b>0.041</b>	-0.5 (0.9)	<b>0.015</b>	-0.2 (0.8)	0.18
Systolic BP (mmHg)	2.7 (11.1)	-1.3 (12.2)	0.20	1.8 (9.7)	0.75	-3.2 (9.4)	<b>0.022</b>
Diastolic BP (mmHg)	0.0 (7.3)	-2.2 (8.5)	0.28	-1.8 (5.3)	0.29	-2.8 (8.3)	0.14
Resting HR (b.p.m.)	-2.7 (6.0)	-5.4 (6.2)	0.11	-6.8 (7.3)	<b>0.020</b>	-6.0 (8.1)	0.07
Fasting glucose (mg/dL)	-0.7 (9.3)	-3.4 (6.2)	0.21	0.2 (7.1)	0.67	-0.7 (7.3)	0.97
Fasting insulin (μIU/mL)	0.06 (2.23)	-0.31 (1.59)	0.47	-0.55 (2.28)	0.29	-0.37 (1.91)	0.39
Total cholesterol	3.6 (25.5)	-10.62 (19.6)	<b>0.023</b>	1.21 (21.5)	0.69	3.09 (28.8)	0.94
Treadmill stress test							
Peak heart rate (b.p.m.)	1.1 (6.0)	-2.5 (7.8)	0.06	-4.8 (5.2)	<b>&lt;0.001</b>	0.2 (6.1)	0.56
VO <sub>2max</sub> (mL/min/kg)	-1.0 (3.1)	2.7 (3.7)	<b>&lt;0.001</b>	2.8 (5.1)	<b>0.001</b>	3.0 (5.9)	<b>0.001</b>
Maximum speed (km/h)	-0.1 (0.6)	1.1 (0.6)	<b>&lt;0.001</b>	1.4 (0.6)	<b>&lt;0.001</b>	0.4 (0.7)	<b>0.002</b>
At 130 b.p.m.	0.2 (0.4)	0.4 (0.6)	<b>0.032</b>	0.6 (0.5)	<b>0.001</b>	0.2 (0.4)	0.82
At 150 b.p.m.	0.2 (0.4)	0.7 (0.7)	<b>&lt;0.001</b>	1.0 (0.5)	<b>&lt;0.001</b>	0.2 (0.4)	0.98
At lactate 2 mmol/L	-0.1 (0.6)	0.8 (0.6)	<b>&lt;0.001</b>	0.5 (0.7)	<b>0.019</b>	-1.0 (0.5)	0.94
RER	-0.04 (0.10)	-0.02 (0.12)	0.45	0.00 (0.16)	0.24	-0.10 (0.13)	<b>0.032</b>

Data are represented as mean (SD).

Significant differences to pre time-point (2-way ANOVA) are in bold face.

b.p.m., beats per minute; BMI, body mass index; BP, blood pressure; Co, control group; HR, heart rate; RER, respiratory exchange ratio.

and E). GEE mixed models confirmed significantly longer telomeres at the post-time point in the AET and IT groups, but not in the control and RT groups. Inclusion of baseline variables such as age, gender, BMI, and baseline VO<sub>2max</sub> did not alleviate this association (Supplementary material online, Table S6). In the second assay, leucocyte telomere length (LTL) was measured in DNA isolates of MNCs by real-time PCR. While the control and RT groups exhibited no significant change after 6 months, LTL significantly increased in both the AET and IT groups (Figure 4A).

### Differential regulation of telomerase activity by endurance and resistance training

The trial revealed no significant change of telomerase activity in the control group and in the resistance exercise group. In contrast, 6 months of AET or high-intensity IT increased telomerase activity by two-fold. The mean relative changes of telomerase activity were control 124 ± 74%, AET 298 ± 334%, IT 264 ± 231%, and RT 169 ± 114% ( $P < 0.05$  for AET and IT vs. control and RT, Figure 4B). Changes of telomerase activity correlated with changes in lymphocyte TL ( $\Delta$  telomerase vs.  $\Delta$  lymphocyte TL:  $R = 0.26$ ,  $P = 0.009$ ).

### Effects of training on cell numbers and telomerase activity in leucocytes sub-populations

In the chronic exercise study, cell numbers of leucocytes, monocytes, lymphocytes, CD3+, CD4+, CD8+, CD19+, CD19+, CD16+/56+, and CD3+HLA-DR+ did not differ between the four study groups and did not differ between the start and the end of the training period. CD34+ haematopoietic progenitor cells were markedly


up-regulated by endurance training and IT, but not in the control or the resistance groups (Supplementary material online, Table S5).

### Insulin-like growth factor-1, alpha-Klotho, irisin, and cortisol concentrations do not correlate with the long-term training effects on telomeres

Insulin-like growth factor-1, alpha-Klotho, and irisin have been proposed to correlate with exercise-induced effects.<sup>16,20,25,26</sup> Alpha-Klotho was discretely up-regulated after 6 months endurance exercise (Supplementary material online, Table S4). No differences were observed for IGF-1 and irisin in the long-term training groups. In the acute exercise study, IGF-1 showed a biphasic response after both running and RT (Supplementary material online, Table S2) as described.<sup>27</sup>

As positive control, cortisol was quantitated in  $n = 10$  participants of the Berlin Marathon.<sup>24</sup> As expected, the physical stress of a marathon markedly increased serum cortisol by 5.5-fold. A 24 h after the marathon, cortisol levels had returned to baseline levels (Figure 5A). In the acute training study, we observed a minor increase of cortisol by 17.8% immediately after running and no change after the resistance protocol. In the prospective long-term exercise study, cortisol serum concentrations were similar between all groups and did not differ before and after the training period (Figure 5B). In the situation of chronic endurance training, the observed chronic changes of telomerase activity appear to be independent of stress hormones.

Nitric oxide (NO) synthases are closely involved in the cellular mechanisms of exercise.<sup>15</sup> Protective signalling effects of running in the vessel wall depend on both endothelial NO synthase and telomerase activity.<sup>13</sup> In blood cells, the inducible isoform of NOS



**Figure 2** Design of the prospective, randomized, and controlled 6 months training study. Individuals enrolled in the study were randomized to the control group (no change of inactive life style) or one of three training groups (three sessions per week, 45 min each): (i) aerobic endurance training (continuous running); (ii) high-intensity interval training (4 × 4 method); (iii) resistance training: circle training on eight strength–endurance training devices. Blood samples were drawn at baseline (pre) and after 6 months (post). \*Discontinuation for orthopaedic reasons was not training-associated. #Plausible stress test was defined as a respiratory exchange ratio of >1.00 in both stress tests.

(iNOS) is predominant. Recent evidence revealed that iNOS is crucial for arterial collateral formation in the context of running.<sup>17</sup> Here, iNOS mRNA expression was up-regulated in both endurance exercise groups, but not with RT (Figure 5C).

### Association of training response with cellular response


We hypothesized that the cellular response could be predicted by changes in physical performance. The mean  $VO_{2max}$  increase between training groups was comparable (Table 2, Supplementary material online, Table S4), however, in individuals with a training effect above the mean of  $\Delta VO_{2max}$  increases, the activity of the telomerase after 6 months training was higher compared with weaker responders (Figure 5D). Changes in TL were not associated with changes of  $VO_{2max}$ .

## Discussion

This is the first prospective, randomized controlled training trial assessing the effects of different training protocols on telomerase

activity and LTL in a primary prevention cohort. The main novel finding of the study is that the activity of the telomerase and TL in circulating MNCs is increased by endurance training and by high-intensity IT but not after RT. These results characterize the cellular ‘anti-aging’ effects of exercise, imply that telomeres adapt to physiological stress and identify differential cellular effects of established training modalities (Take home figure). In middle-aged subjects without a relevant training history, completing a training programme of moderate or intensive endurance exercise led to potent ‘anti-aging effects’ in circulating blood cells, consistent with beneficial effects on cardiovascular health in the long term.

Telomeres cap the ends of all eukaryotic chromosomes and are a prerequisite of genomic stability.<sup>6</sup> During the process of natural aging of somatic cells, TL is one of the major determinants of the cells’ capability to divide and function. The process of telomere attrition plays both a causal as well as a potentiating role in human disease processes and is susceptible to life style factors.<sup>7,10,28</sup> Telomere length in circulating leucocytes has been linked to cardiovascular risk factors and diseases.<sup>6,8–11,29</sup> Physical exercise is a regulator of cellular senescence and TL.<sup>10,12,13,16,18,19</sup> However, previous training studies in




**Figure 3** Endurance but not resistance training induces telomere elongation in lymphocytes and granulocytes. Telomere length was measured in mononuclear cells using the FlowFISH method. Absolute values of telomere length (kbp) were calculated using Southern blot standardization. Individual absolute values of (A) lymphocyte and (B) granulocyte telomere length pre- and post-study time points for each group and means of each group are shown as bold blue bars. AET, aerobic endurance training; IT, interval training; RT, resistance training. (C) FlowFISH FACS plots showing telomere fluorescence intensity peaks for a representative subject of each study group (columns) and pre- vs. post-time points (upper and lower row, respectively). Red peaks are the internal standard (bovine thymocytes) in each sample, blue peaks are lymphocytes, and green peaks are granulocyte measurements. Dotted lines are for visualization of rightward or leftward shifts in the peaks during the study. (D) Mean and SD of relative individual changes of lymphocyte (left panel) and (E) granulocyte (right panel) telomere length for each group.

humans did not provide a conclusive answer with regard to telomere biology because of lack of prospective randomization, a control group, methodological homogeneity, and sufficient duration.<sup>12</sup> To our knowledge, no randomized controlled prospective studies comparing different exercise modalities with respect to cellular senescence have been performed to date.

Data on the acute effects of exercise on TL and telomere-associated proteins after a single or a few bouts of exercise are scarce, especially with regard to RT.<sup>30,31</sup> Here, we present the first data of a cross-over study in healthy young volunteers directly comparing endurance and resistance exercise. A single bout of AET (running) acutely up-regulated MNC telomerase activity. The novel and






**Figure 4** Differential effects of training on telomerase activity and telomere length in circulating mononuclear cells. (A) Leukocyte telomere length was determined in genomic DNA isolates by real-time PCR. Beside telomere (T) DNA, the single (S) copy gene 36b4 was amplified to allow calculation of T/S ratio for each subject at both time points. Data are represented as mean  $\pm$  SD of T/S ratio per group. (B) Telomerase activity was determined by TRAP assay. Data are represented as mean  $\pm$  SD of telomerase activity expressed as human embryonic kidney cell equivalents.

unexpected finding is that circle training on eight strength devices did not induce these cellular effects in the same individuals. Magnet-activated cell sorting with telomerase repeat-amplification protocol (MACS-TRAP) assays revealed increased telomerase activity in CD34<sup>+</sup> and in CD14<sup>+</sup> cells that was observed immediately after endurance exercise and persisted after 24 h. Similarly, up-regulation of telomerase activity was specific for the endurance protocol.

The participants in the training groups exercised three times per week for 26 weeks. The blood draw was performed between 48 h and

7 days after the last training bout. In addition to the comparison of traditional endurance training and RT, high-intensity IT was included. Low-volume high-intensity interval exercise may induce more potent effects on fitness and metabolism, both in healthy individuals as well as patients with heart failure.<sup>5</sup> The IT led to comparable changes in physical fitness compared with AET. These data confirm the recent SAINTEX-CAD study.<sup>32</sup> While acute endurance exercise-induced up-regulation of leucocytes,<sup>33</sup> the cell numbers of leucocytes, monocytes, lymphocytes, CD3<sup>+</sup>, CD4<sup>+</sup>, CD8<sup>+</sup>, CD19<sup>+</sup>, CD19<sup>+</sup>, CD16<sup>+</sup>/56<sup>+</sup>, CD3<sup>+</sup>, and




**Figure 5** Cortisol production in acute extreme exercise vs. chronic exercise and training response. (A) Serum concentrations of the stress hormone cortisol were measured in  $N = 10$  male samples of the Berlin Beat of Running marathon study and compared with (B) cortisol concentrations in 40 samples (10 per experimental group) of the chronic exercise study. (C) mRNA expression of the inducible nitric oxide synthase (real-time PCR). (D) Training response is associated with markers of cellular senescence: responders in the training groups with respect to the change of maximum oxygen uptake ( $\Delta VO_{2max}$ ) after the study compared with baseline were defined as the individuals above the mean of  $\Delta VO_{2max}$  as opposed to 'weak' responders with a  $\Delta VO_{2max}$  below the mean. Comparison of the change in telomerase activity shown as  $\Delta$  human embryonic kidney cell equivalents between subjects below vs. above the mean of the  $\Delta VO_{2max}$ .

HLA-DR+CD3+ did not differ in the chronic exercise study between the four study groups and did not differ between the start and the end of the training period. CD34+ haematopoietic progenitor cells were up-regulated by endurance training and IT as expected,<sup>15,34</sup> but not in the control or the endurance groups. Assessment of stress hormones confirmed marked up-regulation of cortisol after the extreme physical

stress of marathon running.<sup>35</sup> However, the chronic levels of cortisol did not differ between the four groups before and after the training period in the chronic exercise study.

To test whether the increased telomerase activity will persist long term with endurance training, a prospective randomized training study including a control group was conducted. The main finding of



our training study is the differential effects of the three training modalities on TL. In order to account for the methodological challenges of TL measurements in humans, especially with regard to PCR,<sup>36</sup> two established independent methods were used.<sup>8,12,13,22,23</sup> Both assays were carefully controlled (details described in [Supplementary material](#) online, *Methods*). In addition, the findings are supported by the measurements of telomerase activity. While the two endurance-based training protocols markedly increased telomerase activity and TL after 6 months, the strength-based protocol did not.<sup>37</sup> So far, only one prospective study addressed the effects of training on TL.<sup>38</sup> Melk *et al.* trained  $n = 59$  men for 210 minutes per week for 6 months. Consistent with our findings they found an increase in TL and telomerase activity. However, that study was not controlled or randomized, did not test IT or RT and did not report which methodology was used for TL measurement. We would like to highlight the importance of a prospective randomized design, a control group and a sufficient follow-up for the interpretation of a training study.

Endurance training and RT induce a number of differential haemodynamic, metabolic, and/or neurohumoral responses, both acute and chronic.<sup>4,39,40</sup> Precise comparisons between endurance and a resistance exercise protocols are scarce. Our intra-individual comparisons of heart rate during acute AET, intensive IT, and resistance exercise ([Supplementary material online, Figure S8](#)) showed that the mean and the maximum heart rate are higher in the endurance training modalities. This may suggest that, compared with resistance exercise, endurance training may induce a higher rate of (laminar) vascular shear stress, which may, e.g. via NO, potentially contribute to the observed cellular effects.<sup>17</sup> Endothelial NO synthase and telomerase activity have shown to be linked in a signalling pathway mediating

exercise-induced vascular protection.<sup>13</sup> In agreement with this hypothesis, we observed a differential regulation of iNOS which was only up-regulated by the endurance training protocols. From an evolutionary perspective, endurance training and high-intensity IT may mimic advantageous travelling and fight and fly behaviour better than strength training.<sup>19</sup> Our data set the stage for future studies to elucidate the details of these regulatory processes. While acute physical stress leads to up-regulation of stress hormones and leucocytes, these alterations are not observed chronically in the basal state after long-term regular exercising. However, we believe that it is likely that the repeated bouts of stress-induced changes induced by the endurance protocols three times a week lead to a shift towards increased baseline telomerase activity when applied long term. This hypothesis represents a possible explanation for the increase of TL with endurance, but not resistance exercise.

Telomere length has become a widely accepted molecular/cellular biomarker of aging.<sup>6</sup> We and others have demonstrated that changes in blood cell telomeres and their associated factors reflect changes in the vessel wall and the myocardium.<sup>13,14</sup> This may be important for the mechanism of cardiovascular prevention by exercise, because increased telomere erosion is associated with disease incidence and severity.<sup>8,11,41</sup> Importantly, telomere dysfunction by chronic inflammation, as in cardiovascular disease, can induce cellular senescence without telomere shortening.<sup>42</sup> Training responses are characterized by an inter-individual variability. Our study shows that changes in telomerase activity correlate not only with the training modality but also with the changes of the individual performance. Measurements of TL could therefore be a useful indicator of 'biological age' in future intervention studies.

## Limitations

The sample size is smaller than other cardiovascular studies, however, this trial is the largest prospective randomized controlled training study performed to date regarding the cellular effects of exercise that includes randomization to well-defined and supervised 6 months training protocols. The study included a sedentary control group. Adherence is the primary challenge of all randomized interventions that address lifestyle changes, especially exercise. This is reflected by drop-out rates in the literature which are comparable to our study.<sup>43,44</sup> The protocol involved fresh isolation of MNC from peripheral blood after each blood draw. Therefore, incomplete samples in very few patients are expected. Per design, the increase of aerobic capacity was similar between the three training groups. Minor differences of the training intensity are possible but inherent to the respective modalities. The every-day activities of the participants outside the supervised training sessions may have contained elements of the other modalities, but this most likely was also the case in the control group during the 6 months observation period.

## Conclusion

In summary, this randomized controlled training study shows that specific modalities of physical exercise mediate differential effects on regulators of cellular senescence. The activity of telomerase and TL is increased by endurance training and high-intensive IT but not after strength training. The data improve the molecular understanding of the protective effects of exercise in primary prevention and underline the potency of physical training in reducing the impact of age-related diseases. Telomerase activity is a sensitive parameter to measure preventive effects of exercise on the cellular level, both acute and chronic. Therefore, cellular senescence markers could be useful parameters to guide the efficacy of preventive exercise programmes. The effects on clinical outcomes need to be scrutinized in a large prospective trial, taking potential differences of training modalities into consideration.

With regard to training recommendations for the prevention of cellular aging, our data support the ESC's current guideline recommendations that resistance exercise should be complimentary to endurance training rather than a substitute.<sup>2</sup>

## Supplementary material

Supplementary material is available at *European Heart Journal* online.

## Acknowledgements

The authors thank Christian Cassel, Steffi Lieblang-Alff, Simone Bamberger, Claudia Schormann, Ellen Becker, Simone Jäger, and Julia Weber for excellent technical assistance. We thank the staff of the Institut für Sport und Präventivmedizin and the Deutsche Hochschule für Prävention und Gesundheitsmanagement (Saarbrücken, Germany) for screening, testing and training the study subjects in the long-term training study. We are indebted to Prof Stefan Wagenpfeil, Institut für Medizinische Biometrie, Epidemiologie und Medizinische Informatik (Homburg, Germany) for guidance with the statistical revision. We thank Georg Häusler and Matthias Endres, Berlin, for the

samples of the Berlin Marathon participants and the staff of smileBEST fitness (Homburg, Germany) for their support of the acute exercise study. Last but not least, we thank all study subjects for taking part.

## Funding

This work was supported by the Corona Foundation (S199/10060/2014), the Deutsche Forschungsgemeinschaft (SFB TRR 219), Saarland University, Saarland Ministry of the Interior, and Leipzig University, Germany.

**Conflict of interest:** none declared.

## References

- Schuler G, Adams V, Goto Y. Role of exercise in the prevention of cardiovascular disease: results, mechanisms, and new perspectives. *Eur Heart J* 2013;**34**: 1790–1799.
- Piepoli MF, Hoes AW, Agewall S, Albus C, Brotons C, Catapano AL, Cooney MT, Corra U, Cosyns B, Deaton C, Graham I, Hall MS, Hobbs FD, Lochan ML, Lollgen H, Marques-Vidal P, Perk J, Prescott E, Redon J, Richter DJ, Sattar N, Smulders Y, Tiberi M, van der Worp HB, van Dis I, Verschuren WM. 2016 European guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J* 2016;**37**:2315–2381.
- Kraus WE, Bittner V, Appel L, Blair SN, Church T, Despres JP, Franklin BA, Miller TD, Pate RR, Taylor-Piliae RE, Vafiadis DK, Whetsel L. The National Physical Activity Plan: a call to action from the American Heart Association. *Circulation* 2015;**131**:1932–1940.
- Pollock ML, Franklin BA, Balady GJ, Chaitman BL, Fleg JL, Fletcher B, Limacher M, Pina IL, Stein RA, Williams M, Bazzarre T. AHA Science Advisory. Resistance exercise in individuals with and without cardiovascular disease: benefits, rationale, safety, and prescription. *Circulation* 2000;**101**:828–833.
- Weston KS, Wisloff U, Coombes JS. High-intensity interval training in patients with lifestyle-induced cardiometabolic disease: a systematic review and meta-analysis. *Br J Sports Med* 2014;**48**:1227–1234.
- Lopez-Otin C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. *Cell* 2013;**153**:1194–1217.
- Xie Z, Jay KA, Smith DL, Zhang Y, Liu Z, Zheng J, Tian R, Li H, Blackburn EH. Early telomerase inactivation accelerates aging independently of telomere length. *Cell* 2015;**160**:928–939.
- Brouillette SW, Moore JS, McMahon AD, Thompson JR, Ford I, Shepherd J, Packard CJ, Samani NJ, West Of Scotland C, Prevention Study G. Telomere length, risk of coronary heart disease, and statin treatment in the West of Scotland Primary Prevention Study: a nested case-control study. *Lancet* 2007;**369**:107–114.
- Farzaneh-Far R, Cawthon RM, Na B, Browner WS, Schiller NB, Whooley MA. Prognostic value of leukocyte telomere length in patients with stable coronary artery disease: data from the Heart and Soul Study. *Arterioscler Thromb Vasc Biol* 2008;**28**:1379–1384.
- Ornish D, Lin J, Chan JM, Epel E, Kemp C, Weidner G, Marlin R, Frenda SJ, Magbanua MJ, Daubenmier J, Estay I, Hills NK, Chainani-Wu N, Carroll PR, Blackburn EH. Effect of comprehensive lifestyle changes on telomerase activity and telomere length in men with biopsy-proven low-risk prostate cancer: 5-year follow-up of a descriptive pilot study. *Lancet Oncol* 2013;**14**:1112–1120.
- Wong LS, de Boer RA, Samani NJ, van Veldhuisen DJ, van der Harst P. Telomere biology in heart failure. *Eur J Heart Fail* 2008;**10**:1049–1056.
- Mundstock E, Zatti H, Louzada FM, Oliveira SG, Guma FT, Paris MM, Rueda AB, Machado DG, Stein RT, Jones MH, Sarria EE, Barbe-Tuana FM, Mattiello R. Effects of physical activity in telomere length: systematic review and meta-analysis. *Ageing Res Rev* 2015;**22**:72–80.
- Werner C, Fürster T, Widmann T, Pöss J, Roggia C, Hanhoun M, Scharhag J, Büchner N, Meyer T, Kindermann W, Haendeler J, Böhm M, Laufs U. Physical exercise prevents cellular senescence in circulating leukocytes and in the vessel wall. *Circulation* 2009;**120**:2438–2447.
- Wilson WR, Herbert KE, Mistry Y, Stevens SE, Patel HR, Hastings RA, Thompson MM, Williams B. Blood leucocyte telomere DNA content predicts vascular telomere DNA content in humans with and without vascular disease. *Eur Heart J* 2008;**29**:2689–2694.
- Laufs U, Werner N, Link A, Endres M, Wassmann S, Jurgens K, Miche E, Böhm M, Nickenig G. Physical training increases endothelial progenitor cells, inhibits neointima formation, and enhances angiogenesis. *Circulation* 2004;**109**:220–226.
- Werner C, Hanhoun M, Widmann T, Kazakov A, Semenov A, Pöss J, Bauersachs J, Thum T, Pfreundschuh M, Müller P, Haendeler J, Böhm M, Laufs U. Effects of

- physical exercise on myocardial telomere-regulating proteins, survival pathways, and apoptosis. *J Am Coll Cardiol* 2008;**52**:470–482.
17. Schirmer SH, Millenaar DN, Werner C, Schuh L, Degen A, Bettink SI, Lipp P, van Rooijen N, Meyer T, Böhm M, Laufs U. Exercise promotes collateral artery growth mediated by monocytic nitric oxide. *Arterioscler Thromb Vasc Biol* 2015;**35**:1862–1871.
  18. Ludlow AT, Ludlow LW, Roth SM. Do telomeres adapt to physiological stress? Exploring the effect of exercise on telomere length and telomere-related proteins. *Biomed Res Int* 2013;**2013**:1.
  19. Haycock PC, Heydon EE, Kaptoge S, Butterworth AS, Thompson A, Willeit P. Leucocyte telomere length and risk of cardiovascular disease: systematic review and meta-analysis. *BMJ* 2014;**349**:g4227.
  20. Hecksteden A, Wegmann M, Steffen A, Kraushaar J, Morsch A, Ruppenthal S, Kaestner L, Meyer T. Irisin and exercise training in humans—results from a randomized controlled training trial. *BMC Med* 2013;**11**:235.
  21. Kim NW, Piatyszek MA, Prowse KR, Harley CB, West MD, Ho PL, Coviello GM, Wright WE, Weinrich SL, Shay JW. Specific association of human telomerase activity with immortal cells and cancer. *Science* 1994;**266**:2011–2015.
  22. Cawthon RM. Telomere measurement by quantitative PCR. *Nucleic Acids Res* 2002;**30**:e47.
  23. Baerlocher GM, Vulto I, de Jong G, Lansdorp PM. Flow cytometry and FISH to measure the average length of telomeres (flow FISH). *Nat Protoc* 2006;**1**:2365–2376.
  24. Schwarz V, Düsing P, Liman T, Werner C, Herm J, Bachelier K, Krüll M, Brechtel L, Jungehulsing GJ, Haverkamp W, Böhm M, Endres M, Haeusler KG, Laufs U. Marathon running increases circulating endothelial- and thrombocyte-derived microparticles. *Eur J Prev Cardiol* 2018;**25**:317–324.
  25. Bostrom P, Wu J, Jedrychowski MP, Korde A, Ye L, Lo JC, Rasbach KA, Bostrom EA, Choi JH, Long JZ, Kajimura S, Zingaretti MC, Vind BF, Tu H, Cinti S, Hojlund K, Gygi SP, Spiegelman BM. A PGC1- $\alpha$ -dependent myokine that drives brown-fat-like development of white fat and thermogenesis. *Nature* 2012;**481**:463–468.
  26. Matsubara T, Miyaki A, Akazawa N, Choi Y, Ra SG, Tanahashi K, Kumagai H, Oikawa S, Maeda S. Aerobic exercise training increases plasma Klotho levels and reduces arterial stiffness in postmenopausal women. *Am J Physiol Heart Circ Physiol* 2014;**306**:H348–H355.
  27. Eliakim A, Cooper DM, Nemet D. The GH-IGF-I response to typical field sports practices in adolescent athletes: a summary. *Pediatr Exerc Sci* 2014;**26**:428–433.
  28. Bennaceur K, Atwill M, Al Zhrany N, Hoffmann J, Keavney B, Breault D, Richardson G, von Zglinicki T, Saretzki G, Spyridopoulos I. Atorvastatin induces T cell proliferation by a telomerase reverse transcriptase (TERT) mediated mechanism. *Atherosclerosis* 2014;**236**:312–320.
  29. Spyridopoulos I, Hoffmann J, Aicher A, Brummendorf TH, Doerr HW, Zeiher AM, Dimmeler S. Accelerated telomere shortening in leukocyte subpopulations of patients with coronary heart disease: role of cytomegalovirus seropositivity. *Circulation* 2009;**120**:1364–1372.
  30. Chilton WL, Marques FZ, West J, Kannourakis G, Berzins SP, O'Brien BJ, Charchar FJ. Acute exercise leads to regulation of telomere-associated genes and microRNA expression in immune cells. *PLoS One* 2014;**9**:e92088.
  31. Zietzer A, Buschmann EE, Janke D, Li L, Brix M, Meyborg H, Stawowy P, Jungk C, Buschmann I, Hillmeister P. Acute physical exercise and long-term individual shear rate therapy increase telomerase activity in human peripheral blood mononuclear cells. *Acta Physiol* 2017;**220**:251–262.
  32. Pattyn N, Vanhees L, Cornelissen VA, Coeckelberghs E, De Maeyer C, Goetschalckx K, Possemiers N, Wuyts K, Van Craenenbroeck EM, Beckers PJ. The long-term effects of a randomized trial comparing aerobic interval versus continuous training in coronary artery disease patients: 1-year data from the SAINTEX-CAD study. *Eur J Prev Cardiol* 2016;**23**:1154–1164.
  33. Witard OC, Turner JE, Jackman SR, Tipton KD, Jeukendrup AE, Kies AK, Bosch JA. High-intensity training reduces CD8<sup>+</sup> T-cell redistribution in response to exercise. *Med Sci Sports Exerc* 2012;**44**:1689–1697.
  34. Walthert C, Gaede L, Adams V, Gelbrich G, Leichtle A, Erbs S, Sonnabend M, Fikenzler K, Korner A, Kiess W, Bruegel M, Thiery J, Schuler G. Effect of increased exercise in school children on physical fitness and endothelial progenitor cells: a prospective randomized trial. *Circulation* 2009;**120**:2251–2259.
  35. Dimitrov S, Benedict C, Heutling D, Westermann J, Born J, Lange T. Cortisol and epinephrine control opposing circadian rhythms in T cell subsets. *Blood* 2009;**113**:5134–5143.
  36. Bateson M, Nettle D. The telomere lengthening conundrum—it could be biology. *Aging Cell* 2017;**16**:312–319.
  37. Borghini A, Giardini G, Tonacci A, Mastorci F, Mercuri A, Mrakic-Spota S, Spota SM, Moretti S, Andreassi MG, Pratali L. Chronic and acute effects of endurance training on telomere length. *Mutagenesis* 2015;**30**:711–716.
  38. Melk A, Tegtbur U, Hilfiker-Kleiner D, Eberhard J, Saretzki G, Eulert C, Kerling A, Nelius AK, Homme M, Strunk D, Berliner D, Rontgen P, Kuck M, Bauersachs J, Hilfiker A, Haverich A, Bara C, Stiesch M. Improvement of biological age by physical activity. *Int J Cardiol* 2014;**176**:1187–1189.
  39. Fowler RM, Maiorana AJ, Jenkins SC, Gain KR, O'Driscoll G, Gabbay E. A comparison of the acute haemodynamic response to aerobic and resistance exercise in subjects with exercise-induced pulmonary arterial hypertension. *Eur J Prev Cardiol* 2013;**20**:605–612.
  40. Lavie CJ, Arena R, Swift DL, Johannsen NM, Sui X, Lee DC, Earnest CP, Church TS, O'Keefe JH, Milani RV, Blair SN. Exercise and the cardiovascular system: clinical science and cardiovascular outcomes. *Circ Res* 2015;**117**:207–219.
  41. Samani NJ, Boulton R, Butler R, Thompson JR, Goodall AH. Telomere shortening in atherosclerosis. *Lancet* 2001;**358**:472–473.
  42. Jurk D, Wilson C, Passos JF, Oakley F, Correia-Melo C, Greaves L, Saretzki G, Fox C, Lawless C, Anderson R, Hewitt G, Pender SL, Fullard N, Nelson G, Mann J, van de Sluis B, Mann DA, von Zglinicki T. Chronic inflammation induces telomere dysfunction and accelerates ageing in mice. *Nat Commun* 2014;**2**:4172.
  43. Miller FL, O'Connor DP, Herring MP, Sailors MH, Jackson AS, Dishman RK, Bray MS. Exercise dose, exercise adherence, and associated health outcomes in the TIGER study. *Med Sci Sports Exerc* 2014;**46**:69–75.
  44. McDermott MM, Ades P, Guralnik JM, Dyer A, Ferrucci L, Liu K, Nelson M, Lloyd-Jones D, Van Horn L, Garside D, Kibbe M, Domanchuk K, Stein JH, Liao Y, Tao H, Green D, Pearce WH, Schneider JR, McPherson D, Laing ST, McCarthy WJ, Shroff A, Criqui MH. Treadmill exercise and resistance training in patients with peripheral arterial disease with and without intermittent claudication: a randomized controlled trial. *JAMA* 2009;**301**:165–174.